#### OHSU DIGESTIVE HEALTH

# Preparing for your upper GI endoscopy with Bravo PH testing

This guide will help you and your family prepare for your procedure. If you have any questions, please call **503-494-4373**. For questions after 5 p.m., call 503-494-8311 and ask for the GI doctor on call.

### Once you are scheduled



- □ Download directions to your check-in location at <a href="https://www.ohsu.edu/digestive-health/preparing-your-endoscopy-procedure">https://www.ohsu.edu/digestive-health/preparing-your-endoscopy-procedure</a>
- ☐ Ask a friend or family member to come with you to your procedure. You should have 1 healthy adult (age 18+) to take you home.
- ☐ Let us know if you have an allergy to nickel or if your skin reacts badly to some types of jewelry.
- ☐ You may need to make major changes to your medications for this procedure. Please review the **medication instructions** on page 2. If you need more instructions on any of your medications, please contact the doctor who prescribed them.
  - ➤ If you get heartburn because you had to stop your medications for the procedure, you can take Tums up until 24 hours before your procedure.





If you take	Do this
Proton Pump Inhibitors - PPIs	
(Omeprazole, Prilosec,	procedure.
Nexium, Protonix, etc.)	
Antacids (Zantac, Tagamet,	STOP taking 3 days before your
Pepcid)	procedure.
Suboxone	Call us at 503-494-4373.
Diabetes medicine by mouth	Do NOT take the night before or the
(Metformin, Glipizide)	morning of your procedure.
Insulin	Take ½ your dose the night before
	and NO insulin the day of your
	procedure. Or, follow the instructions
	from the OHSU Preoperative
	Medicine Clinic. Your blood sugar
<u> </u>	could be lower from no solid food.
Benzodiazepines	STOP taking 2 days (48 hours) before
(clonazepam, Klonipin,	your procedure.
diazepam, Valium, Iorazepam,	
Ativan, midazolam, Versed,	
temazepam, Restoril) Bentyl (dicyclomine)	STOP taking 24 hours before.
• • • • • • • • • • • • • • • • • • • •	
Blood thinners*	Talk to the doctor who prescribed
	your medication to talk about the
* Coumadin or Warfarin pills	guidelines below. STOP taking 5 days before your
Countain of Wartain pins	procedure if OK with your prescribing
	doctor. Call us if you CANNOT stop.
* Levenox or Enoxaparin	STOP taking 24 hours before your
injections	procedure if OK with your prescribing
	doctor. Call us if you CANNOT stop.
* Novel anticoagulant pills	STOP taking 1 day before your
(Paradaxa, Dabigatran,	procedure if OK with your prescribing
Eliquis, Apixaban, Xarelto,	doctor. Call us if you CANNOT stop.
Riveroxaban, Edoxaban,	
Savaysa)	
* Anti-platelets (Plavix,	STOP taking 7 days before your
Clopidogrel, Prasugrel,	procedure if OK with your prescribing
Effient)	doctor. Call us if you CANNOT stop.

☑ Keep taking all of your other prescribed medicines, including aspirin, pain medications, blood pressure medications like Lisinopril, water pills (diuretics like Lasix), Non-steroidal anti-inflammatory drugs (like Advil, Motrin, Aleve).



## 1 day before your procedure



- You may eat normally until midnight before your procedure.
- **NO solid food after midnight.**

## The day of your procedure



• Only drink clear liquids (liquids you can see through when you hold it up to the light) until 2 hours before your arrival time. Do NOT eat any solid food. Follow this schedule no matter what time of day your procedure is.

#### **☑ DO**:

- Drink clear (**not red or purple**) liquids like:
  - ☑ Clear broth (chicken, beef or vegetable)
  - ☑ Coffee or tea (no milk, but sugar is OK)
  - ☑ Carbonated soft drinks
  - ☑ Clear juice like apple and white grape (no pulp)
  - ☑ Clear sports drinks like Gatorade and Powerade (no red or purple)
  - ☑ Popsicles and Jell-O (no red or purple)

#### $\bigcirc$ **DON'T**:

- Drink liquids that are purple or red.
- O Drink milk, orange juice, tomato juice, broth that has solids, or solid food.
- > 2 hours before your arrival time, stop drinking anything.
  - You can have clear liquids up until 2 hours before your arrival time.
  - o You may take your medicine with a sip of water.



## After your procedure



- You will carry a wireless monitor that can attach to your belt for 2 days.
- We will schedule you for an appointment 2 days later to return the monitor.
- You must bring it back in 2 days so we can download the data. You MAY NOT mail it back but you may have someone else bring it back for you.

